§ 520.90

such as wounds, abscesses and cellulitis/dermatitis due to susceptible strains of beta-lactamase (penicillinase) producing S. aureus, nonbeta-lactamase S. aureus, Staphylococcus spp., Streptococcus spp., E. coli, Pasteurella multocida, and Pasteurella spp.

(iii) Limitations. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37320, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995; 63 FR 13121, Mar. 18, 1998; 79 FR 28817, May 20, 2014]

§520.90 Ampicillin oral dosage forms.

§520.90a [Reserved]

§520.90b Ampicillin tablets.

- (a) Specifications. Each tablet contains ampicillin trihydrate equivalent to 50 or 100 milligrams of ampicillin.
- (b) Sponsor. See No. 054771 in $\S 510.600(c)$ of this chapter.
- (c) Conditions of use in dogs—(1) Amount. 5 milligrams per pound of body weight, at 8-hour intervals, 1 to 2 hours prior to feeding, to be continued 36 to 48 hours after all symptoms have subsided. If no improvement is seen within 5 days, stop treatment, reevaluate diagnosis, and change therapy.
- (2) Indications for use. Oral treatment of infections caused by susceptible organisms as follows: Upper respiratory infections, tonsillitis, and bronchitis due to Streptococcus spp., Staphylococcus spp., Escherichia coli, Proteus mirabilis, and Pasteurella spp., urinary tract infections (cystitis) due to Streptococcus spp., Staphylococcus spp., E., coli, P. mirabilis, and Enterococcus spp.; gastrointestinal infections due to Staphylococcus spp., Streptococcus spp.. Enterococcus spp., and E. coli.; infections associated with abscesses, lacerations, and wounds caused by Staphylococcus spp., and Streptococcus spp.
- (3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37321, Aug. 18, 1992, as amended at 60 FR 55659, Nov. 2, 1995; 79 FR 28818, May 20, 2014]

§520.90c Ampicillin capsules.

(a) Specifications. Each capsule contains ampicillin trihydrate equivalent

- to 125, 250, or 500 milligrams of ampicillin.
- (b) Sponsor. See No. 054771 in $\S510.600$ (c) of this chapter.
- (c) Conditions of use—(1) Dogs—(i) Amount. 5 to 10 milligrams per pound of body weight two or three times daily. In severe or acute conditions, 10 milligrams per pound of body weight, three times daily. Administer 1 to 2 hours prior to feeding.
- (ii) Indications for use. Treatment against strains of gram-negative and gram-positive organisms sensitive to ampicillin and associated with respiratory tract infections (tracheobronchitis and tonsillitis); urinary tract infections (cystitis); bacterial gastroenteritis; generalized infections (septicemia) associated with abscesses, lacerations, and wounds; and bacterial dermatitis.
- (iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Cats—(i) Amount. 10 to 30 milligrams per pound of body weight or three times daily. Administer 1 to 2 hours prior to feeding.
- (ii) Indications for use. Treatment against strains of gram-negative and gram-positive organisms sensitive to ampicillin and associated with respiratory tract infections (bacterial pneumonia); urinary tract infections (cystitis); and generalized infections (septicemia) associated with abscesses, lacerations, and wounds.
- (iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37321, Aug. 18, 1992, as amended at 58 FR 61016, Nov. 19, 1993; 79 FR 28818, May 20, 2014]

§ 520.90d Ampicillin for oral suspension.

- (a) Specifications. When reconstituted as directed, each milliliter contains ampicillin trihydrate equivalent to 25 milligrams of ampicillin.
- (b) Sponsor. See No. 055529 in \$510.600(c) of this chapter.
- (c) Conditions of use—(1) Dogs—(i) Amount. Administer to 10 milligrams per pound of body weight orally, 2 or 3 times daily, 1 to 2 hours prior to feeding. In severe or acute conditions, 10 milligrams per pound of body weight 3